(12) UK Patent Application (19) GB (11) 2 367 493 (13) A

(43) Date of A Publication 10.04.2002

- (21) Application No 0024590.2
- (22) Date of Filing 07.10.2000
- (71) Applicant(s)

Growing Success Organics Limited (Incorporated in the United Kingdom) 1-3 Hill Top Business Park, Devizes Road, SALISBURY, Wiltshire, SP3 4UF, United Kingdom

(72) Inventor(s)

Stanley Edward John Mernagh Alan McClymont

(74) Agent and/or Address for Service

Urquhart-Dykes & Lord 1 Richfield Place, Richfield Avenue, READING, RG1 8EQ, United Kingdom (51) INT CL⁷
A01N 25/30

- (52) UK CL (Edition T)

 A5E EF E300 E303 E318

 U1S S1097 S1102 S1277 S1279 S1290 S1303
- (58) Field of Search
 UK CL (Edition S) **A5E EF**INT CL⁷ **A01N 25/30**Online: EPODOC, JAPIO, WPI

(54) Abstract Title

Composition for deterring rodents comprising a capsicum oleoresin, a polyoxyethylene sorbitan fatty acid ester & a fatty acid triglyceride

- (57) A rodent-deterrent composition comprises:
 - (a) a capsicum oleoresin,
 - (b) a polyoxyethylene sorbitan fatty acid ester,
 - (c) a fatty acid triglyceride, and
 - (d) water

and is in the form of an oil-in-water emulsion. It gives good overall coverage of treated areas.

The capsicum oleoresin may be obtained by extraction of the fruits of Capsicum annum L. or Capsicum frutescens L. The polyoxyethylene sorbitan fatty acid ester is preferably a sorbitan monopoleate or trioleate ester or a sorbitan monopalmitate ester, with about 20 units ethylene oxide. The fatty acid triglyceride is preferably a C₈₋₁₀ saturated fatty acid triglyceride.

A bird feed (eg one comprising seeds, nuts or a fat ball) may be rendered deterrent to rodents by applying the above composition to the feed.

The two surfactants present in the composition are of food grade and are fully biodegradable.

COMPOSITION AND METHOD FOR DETERRING RODENTS

This invention relates to a method for the deterrence of rodents, to a composition for this purpose, and to an animal food treated with such a composition. More particularly, it relates to a method, a composition, and

animal food intended to repel the grey squirrel (Sciurus carolinensis).

Although the grey squirrel was introduced into Britain more than a century ago, it is only in the last two or three decades that it has become so widespread through large areas of the country as to become a major pest, and almost entirely to replace the native red squirrel (Sciurus vulgaris).

The problem of this infestation has become more prevalent because of the increasing practice of putting out food, such as nuts and seeds, intended to attract wild birds to gardens. The grey squirrel is a very efficient forager, and has become very efficient at finding out how to take food from bird feeders. There have even been television programmes displaying their ingenuity and capacity for learning how to circumvent quite complicated barriers.

Although they are not unattractive to the sight, they can constitute a health and economic hazard. As rodents, they are vehicles for the transmission of a variety of bacterial and viral infections, which may be transmitted to humans or domestic animals which come into contact with their urine or faeces. They may also be hosts to

5

10

15

ticks or fleas which may then be transferred to domestic pets. They can, moreover cause severe structural damage to buildings if they should gain access to roof spaces. In addition, they can very readily multiply domestic expenditure on bird seed and nuts.

Various methods have been adopted to deter these pests. For instance there are various barrier systems and traps, but these have been found to be costly and to have only 10 limited effectiveness. Chemical methods have also been proposed, but these also have a number of practical disadvantages. For instance, one method that has been employed is to sprinkle finely ground black pepper or capsicum seed around areas where squirrels enter or 15 congregate. Although this can be effective, in that · squirrels are repelled, there is the disadvantage that long term deterrence has proved difficult for practical reasons. Powders are not easily applied to areas that are exposed, not flat, or inaccessible. For instance, powders 20 cannot be applied to vertical or steeply-sloping areas, such as walls or roofs, or to bird food hanging in vertical nets or cages. Even when applied to horizontal surfaces, such as bird tables, the tops of walls or fences, the roofs of sheds etc., long term protection is 25 difficult or impossible, because even slight breezes will disperse the powder, and render it ineffective. There is therefore a need for a simple means of deterring squirrels or other rodents which is effective and long lasting, and which imposes no health hazards to humans, 30 domestic pets or to bird life. The present invention provides a simple means of meeting these requirements.

In one embodiment the present invention provides a rodent-deterrent composition which comprises:

- (a) a capsicum oleoresin,
- (b) a polyoxyethylene sorbitan fatty acid ester,
- (c) a fatty acid triglyceride, and
- (d) water.

5

Preferably the composition according to the invention comprises:

- (a) from 0.10 to 10.00 %w/w of the capsicum oleoresin,
- 10 (b) from 0.15 to 15.00 %w/w of the polyoxyethylene sorbitan fatty acid ester,
 - (c) from 0.15 to 15.00 %w/w of the fatty acid triglyceride, and $\ensuremath{\mbox{\sc triglyceride}}$
 - (d) from 60.00 to 99.60 %w/w of water.

15

20

· More preferably, the composition comprises:

- (a) from 0.10 to 0.50 %w/w of the capsicum oleoresin,
- (b) from 0.15 to 0.75 %w/w of the polyoxyethylene sorbitan fatty acid ester,
 - (c) from 0.15 to 0.75 %w/w of the fatty acid triglyceride, and
 - (d) from 98.00 to 99.60 %w/w of water.
- 25 The capsicum oleoresin is obtained by extraction in known manner of the fruits of Capsicum annum L. or Capsicum frutescens L. or other Capsicum species.
- Suitable polyoxyethylene sorbitan fatty acid esters include oleic acid esters and palmitic acid esters. Examples of suitable commercially available esters include:

Surfacare T20: Sorbitan trioleate ester with 20 units

(moles) ethylene oxide, (EO)

Crillet 4: Sorbitan monooleaate ester with 20 units

ΕO

5 Tween 4: Sorbitan monopalmitate ester with 20 units

ΕO

Tween 85: Sorbitan trioleate ester with 20 units EO

According to another embodiment, the invention provides a 10 method of rendering a bird feed, such as seeds, nuts or a fat ball, deterrent to rodents which comprises applying a composition as described above.

A suitable commercially available saturated fatty acid triglyceride is Surfac MCTG, a C_{θ} - C_{10} saturated fatty acid triglyceride.

The compositions according to the invention are oil-in-water emulsions which have moderate phase stability and a large droplet size, giving good overall coverage of treated areas. They combine non-toxicity, due to the use of low levels of the capsicum oil and of two surfactants, which are of food grade and are fully biodegradable. They are thus able to combine convenience and effectiveness in use with a high initial activity and low environmental impact.

In the formation of the emulsions, use is made of two surfactants, component (b) acting as the primary

emulsifier, while component (c) acts as secondary emulsifier. It is found that the combination of the two surfactants provides the most acceptable emulsion characteristics.

20

It is found that the compositions according to the invention are effective in deterring rodents, such as grey squirrels, while being palatable to garden birds.

5 If desired, other ingredients, such as food grade dyes, may be incorporated into the compositions for special purposes.

CLAIMS

- A rodent-deterrent composition which comprises:
 - (a) a capsicum oleoresin,
 - (b) a polyoxyethylene sorbitan fatty acid ester,
 - (c) a fatty acid triglyceride, and
 - (d) water.
- 2 A composition according to claim 1 which comprises:
- (a) from 0.10 to 10.00 %w/w of the capsicum oleoresin,
- (b) from 0.15 to 15.00 %w/w of the polyoxyethylene sorbitan fatty acid ester,
- (c) from 0.15 to 15.00 %w/w of the fatty acid triglyceride, and
 - (d) from 60.00 to 99.60 %w/w of water.
 - 3 A composition according to claim 1 or 2 which comprises:
 - (a) from 0.10 to 0.50 %w/w of the capsicum oleoresin,
 - (b) from 0.15 to 0.75 %w/w of the polyoxyethylene sorbitan fatty acid ester,
 - (c) from 0.15 to 0.75 $\mbox{\ensuremath{\$w/w}}$ of the fatty acid triglyceride, and
 - (d) from 98.00 to 99.60 %w/w of water.
 - A composition according to any of claims 1 to 3 wherein the capsicum oleoresin is obtained by extraction of the fruits of Capsicum annum L. or Capsicum frutescens L.
 - 5 A composition according to any one of the preceding

claims wherein the polyoxyethylene sorbitan fatty acid ester is an oleic acid ester or a palmitic acid ester.

- A composition according to claim 5 wherein the polyoxyethylene sorbitan fatty acid ester is a sorbitan monooleate or trioleate ester or a sorbitan monopalmitate ester, with about 20 units ethylene oxide.
- 7 A composition according to any one of the preceding claims wherein the fatty acid triglyceride is a C_8 C_{10} saturated fatty acid triglyceride.
- 8 A method of rendering a bird feed deterrent to rodents which comprises applying a composition according to any of claims 1 to 7 to the feed.
- 9 A method according to claim 8 wherein the bird feed comprises seeds, nuts or a fat ball.







Application No: Claims searched:

GB 0024590.2

1-9

Examiner: Date of search:

Stephen Quick 4 April 2001

Patents Act 1977 Search Report under Section 17

Databases searched:

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:

UK Cl (Ed.S): A5E (EF)

Int Cl (Ed.7): A01N 25/30

Other:

Online: EPODOC, JAPIO, WPI

Documents considered to be relevant:

Category	Identity of document and relevant passage	Relevant to claims
A	US 5607715 A (PROCTER & GAMBLE), see columns 2 (lines 14-22 & 28-31) & 4 (lines 54-57)	-
A	WPI Abstract Accession No 1997-276666/25 & JP 090100206 A 15.04.1997, see abstract (rat repellent comprising capsicum oil)	-

& Member of the same patent family

- Document indicating technological background and/or state of the art.
- P Document published on or after the declared priority date but before the filing date of this invention.
- E Patent document published on or after, but with priority date earlier than, the filing date of this application.

X Document indicating lack of novelty or inventive step
 Y Document indicating lack of inventive step if combined with one or more other documents of same category.